

Ultraviolet irradiation doses for coronavirus inactivation – review and analysis of coronavirus photoinactivation studies

Ultraviolette Bestrahlungsdosen für die Inaktivierung von Coronaviren – Review und Analyse von Coronavirusinaktivierungsstudien

Abstract

Background: To slow the increasing global spread of the SARS-CoV-2 virus, appropriate disinfection techniques are required. Ultraviolet radiation (UV) has a well-known antiviral effect, but measurements on the radiation dose necessary to inactivate SARS-CoV-2 have not been published so far.

Methods: Coronavirus inactivation experiments with ultraviolet light performed in the past were evaluated to determine the UV radiation dose required for a 90% virus reduction. This analysis is based on the fact that all coronaviruses have a similar structure and similar RNA strand length.

Results: The available data reveals large variations, which are apparently not caused by the coronaviruses but by the experimental conditions selected. If these are excluded as far as possible, it appears that coronaviruses are very UV sensitive. The upper limit determined for the log-reduction dose (90% reduction) is approximately 10.6 mJ/cm² (median), while the true value is probably only 3.7 mJ/cm² (median).

Conclusion: Since coronaviruses do not differ structurally to any great extent, the SARS-CoV-2 virus – as well as possible future mutations – will very likely be highly UV sensitive, so that common UV disinfection procedures will inactivate the new SARS-CoV-2 virus without any further modification.

Keywords: coronavirus, SARS-CoV, SARS-CoV-2, MERS-CoV, ultraviolet, UVC, irradiation, inactivation, disinfection, Covid-19

Zusammenfassung

Hintergrund: Um die weltweite Ausbreitung des SARS-CoV-2 Virus zu verlangsamen, werden geeignete Desinfektionstechniken benötigt. Ultraviolette Strahlung (UV) hat bekanntlich eine antivirale Wirkung, aber Messungen zu benötigten Bestrahlungsdosen für die Inaktivierung von SARS-CoV-2 sind bisher nicht veröffentlicht worden.

Material und Methoden: Coronavirusinaktivierungsexperimente, die in der Vergangenheit durchgeführt wurden, werden herangezogen, um die UV-Bestrahlungsdosis für eine 90%ige Virusreduktion zu ermitteln. Die durchgeführte Analyse nutzt dabei die Tatsache, dass alle Coronaviren eine ähnliche Struktur und eine vergleichbare RNA-Länge aufweisen.

Ergebnisse: Die verfügbaren Daten weisen große Variationen auf, die durch unterschiedliche experimentelle Bedingungen zu erklären sind. Wenn extremere Versuchsbedingungen ausgeschlossen werden, zeigt sich, dass Coronaviren sehr UV-empfindlich sind. Der ermittelte obere Grenzwert für die log-Reduktionsdosis (90% Reduktion) beträgt ungefähr 10.6 mJ/cm² (Median), während der wahre Wert vermutlich nur etwa 3.7 mJ/cm² (Median) beträgt.

Martin Heßling¹
Katharina Hönes¹
Petra Vatter¹
Christian Lingenfelder²

1 Institute of Medical Engineering and Mechatronics, Ulm University of Applied Sciences, Ulm, Germany

2 Pharpur GmbH, Königsbrunn, Germany

Schlussfolgerung: Da sich Coronaviren in ihrer Struktur nicht stark unterscheiden, wird auch das neue SARS-CoV-2 Virus – und mögliche zukünftige Mutationen – sehr UV-empfindlich sein, vermutlich sogar so UV-empfindlich, dass etablierte UV-Desinfektionsverfahren das neue SARS-CoV-2 Virus ohne zusätzliche Modifikationen effizient inaktivieren können.

Schlüsselwörter: Coronavirus, SARS-CoV, SARS-CoV-2, MERS-CoV, ultraviolette Strahlung, UVC, Bestrahlung, Inaktivierung, Desinfektion, Covid-19

Introduction

The newest coronavirus disease COVID-19 is a highly transmittable and pathogenic viral infection caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which emerged in Wuhan, China and spread around the world. So far, more than 3.8 million infections have led to more than 265,000 fatalities worldwide [1].

The responsible pathogen, SARS-CoV-2, is an enveloped single-stranded RNA virus and member of the family coronaviridae of the order nidovirales. All viruses of this family exhibit very similar features. They have a spheroid shape with a diameter of about 100–150 nm, are covered with spike proteins on the outside, and have an RNA strand length of 27–32 kb on the inside [2]. Four of these coronaviruses are known to be the causative agents of common colds (HCoV-229E, HCoV-NL63, HCoV-OC43 and HCoV-HKU1), which usually only result in milder infections, but the coronaviruses MERS-CoV (MERS: Middle East Respiratory Syndrome) SARS-CoV and SARS-CoV-2 have claimed many lives [3].

To stop the spread of SARS-CoV-2, several measures such as containment, social distancing and wearing face masks have been taken. Among other steps, hygiene procedures have been intensified. The employment of liquid disinfectants is one procedure that has been successful against the 6 older coronaviruses [4]. Thermal disinfection has also been proven to be quite effective, even at temperatures as low as 60 °C to 80 °C [4].

Radiation disinfection, especially ultraviolet (UV) radiation, is another well-known inactivation approach for all known microorganisms and viruses that offers some advantages over liquid disinfectants and heat sterilization. It can be performed automatically and employed to disinfect surfaces, liquids, air and rooms, and it is also very energy-efficient.

The ultraviolet spectrum is divided in 4 sections: Radiation with a wavelength between 100 and 200 nm is called vacuum ultraviolet radiation (VUV). It is usually not applied for disinfection purposes because it is strongly absorbed by air. The better-known ultraviolet ranges are UVC, UVB, and UVA with spectral ranges of 200–280 nm, 280–315 nm, and 315–380 nm, respectively. Among these last three UV ranges, UVC is the one with the strongest antimicrobial/antiviral properties [5], [6]. For RNA viruses, the main inactivation mechanism is illustrated in Figure 1A. UV radiation is absorbed by the RNA,

which leads to the formation of pyrimidine dimers, e.g., uracil dimers [6], [7].

The most common UVC light sources for many decades now have been mercury discharge lamps, especially low-pressure mercury vapor lamps, with a strong emission peak at 254 nm, which is near the RNA absorption maximum at about 260 nm, as depicted in Figure 1B.

Although it is known that this kind of UVC radiation has an inactivating effect on all microorganisms and viruses, all pathogens require different UVC irradiation doses for successful inactivation. For instance, the rotavirus requires about 25 mJ/cm² of 254 nm UVC radiation from a mercury discharge lamp for a 3-log reduction, but for adenovirus (Type 40), the value is approximately 6 times higher (140 mJ/cm²) [5], [8].

To answer the important question regarding SARS-CoV-2 and other coronaviruses as to which irradiation doses are needed for inactivation, the existing coronavirus photoinactivation results of the last 60 years have been reviewed and analyzed in this study.

Materials and methods

Google Scholar and Pubmed were searched for different combinations of the following terms: coronavirus, inactivation, photoinactivation, disinfection, ultraviolet, radiation, and light. In addition, a number of individual viruses that belong to the families of coronaviruses although the term “coronavirus” does not occur in their name were also searched for, such as porcine epidemic diarrhea virus, transmissible gastroenteritis virus, feline infectious peritonitis virus, mouse hepatitis virus, sialodacryoadenitis virus, hemagglutinating encephalomyelitis virus, and infectious bronchitis virus.

The retrieved sources were evaluated according to the type of sample irradiated (aerosol, surface, or liquid), the type of light source (including emission peak wavelength), the inactivation effect achieved, and the applied irradiation dose.

If information on disinfection results for different irradiation doses could be found in a single article, those describing a virus reduction by approx. 3–4 log levels were selected. Results that were only displayed as illustrations without the corresponding values in the text or tables were read from previously enlarged figures.

If the necessary information was incomplete, e.g., because of missing irradiation details, the irradiation was




Figure 1: A) Scheme of UV-RNA-damaging mechanism by dimer formation. B) Relative absorption spectra of RNA, relative emission spectrum of a low-pressure mercury vapor lamp and transmission of a typical (Eagle) cell culture medium.

estimated by available lamp information, provided that lamp type and distance were given or by other available means. Publications in which radiation was combined with photosensitizers or other chemical or biochemical agents were excluded. Subsequently, these data were employed to calculate the log-reduction dose, i.e., the irradiation dose required for a 90% virus reduction.

Results and discussion

About 30 publications of scientific investigations regarding photoinactivation of coronaviruses were identified. This included studies on CoV, SARS-CoV or MERS-CoV. An overview of the results is presented by sample condition and wavelength in Table 1.

Almost all experiments were performed with mercury vapor lamps, with a peak emission at 254 nm (UVC), which is near the RNA absorption peak in Figure 1. Individual investigations were performed with peak wavelengths at 222 nm (UVC), or 365 nm (UVA), or even with daylight. In most studies, the authors did not intend to investigate the log-reduction doses of coronaviruses, but rather virus inactivation in various applications. In all experiments and for all coronaviruses, a successful virus inactivation was observed. However, because of the different objectives of the studies, the experimental conditions to determine the specific log-reduction doses were often difficult to identify. In many cases, information important for the present study's analysis was missing.

To evaluate photoinactivation results, the basic inputs were the virus reduction and the applied irradiation dose. Not all authors provided the applied irradiation dose, but at least for some studies this value could be calculated as the product of irradiation duration and irradiation intensity, or it could be estimated. For some investigations, it was impossible to quantify the disinfection success exactly; in these cases, the values were estimated based on the information given in Table 1. However, for some

studies, it was even impossible to roughly estimate the log-reduction dose.

The calculated and estimated results, given as log-reduction doses, exhibit extreme variability, even within the 254 nm results, ranging from 0.6 mJ/cm² (bovine corona virus [9]) to 11,754 mJ/cm² (SARS (CoV Urbani) [10]). Even the differences between the SARS-CoV strains were above two orders of magnitude concerning the necessary dose.

Possible reasons for this observation might be biological and biochemical differences between the coronavirus strains. However, comparing the experimental details, two other potential dominant factors attract attention:

1. The necessary irradiation doses are lower for viruses on surfaces, aerosols and pure salt solutions. When irradiation experiments were performed with the virus in different solutions, it is important to bear in mind that the solutions contain organic materials, e.g., blood products or residue from cell culture medium. These solutions exhibit very high absorption of the applied UVC radiation, resulting in much lower irradiances for viruses that are deeper inside the sample. This effect is clearly illustrated in Figure 1B (culture medium transmission) and by the results of Terpstra et al. [11]. Those authors were aware of the absorption of their samples and presented results with 10%- and 30%-virus-containing plasma within the irradiated samples. Although it was the same coronavirus (transmissible gastroenteritis virus (TGEV)) and the same experimental setup, the results differ by a factor of 3.1.

Most authors did not measure the UVC absorption properties of their biological materials because it was of no importance for their research task; thus, it is almost impossible to extract the role of the absorption in the calculation of the necessary irradiation doses for a 90% virus reduction. In consequence, the lower values for the log-reduction doses, mostly from viruses in salt solutions, surfaces or aerosols, might be a more realistic approach to determine the true virus log-reduction dose.

Table 1: Overview of published and analyzed coronavirus photoinactivation investigations sorted by sample condition

Virus	Irradiation wavelength [nm]	Log-reduction dose [mJ/cm ²]	Sample condition	Remark	Reference
Human Coronavirus (HCoV-229E)	222	0.56	aerosol	values calculated by authors	[17]
Human Coronavirus (HCoV-OC43)	222	0.39	aerosol	values calculated by authors	[17]
Murine Coronavirus (MHV)	254	0.66	aerosol	values calculated by Table 1	[18]
Middle East Respiratory Syndrome Coronavirus (MERS-CoV)	254	successful inactivation	droplets	no dose information	[19]
Infectious Bronchitis Coronavirus (IBV)	254	13.84	surface	values taken from Figure 2	[20]
Murine Coronavirus (MHV A59)	254	successful inactivation	surface	no dose information	[19]
Transmissible Gastroenteritis Virus (TGEV)	254	successful inactivation	surface	no dose information	[21]
Transmissible Gastroenteritis Virus (TGEV)	daylight	successful inactivation	surface	no dose information	[21]
Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV)	254	successful inactivation	liquid (salt solution)	no quantification possible	[22]
Berne Torovirus (ToV P 138/72)	254	1.25*	liquid (very thin layer of cell culture medium with fetal bovine serum)	torovirus: no member of coronavirinae (different spatial structure), data not included in analysis	[14]
Canine Coronavirus (CCoV I-71)	254	10.55	liquid (thin layer of cell culture medium with fetal bovine serum)	missing irradiation dose calculated by lamp data and irradiation time	[23]
Murine Coronavirus (MHV 2)	254	1.54	liquid (very thin layer of cell culture medium with fetal bovine serum)	probably not highly absorbed	[24]
Murine Coronavirus (MHV 2)	254	8.67	liquid (thin layer of cell culture medium with fetal bovine serum)	missing irradiation dose calculated by lamp data and irradiation time	[23]
Murine Coronavirus (MHV N)	254	12.1	liquid (thin layer of cell culture medium with fetal bovine serum)	missing irradiation dose calculated by lamp data and irradiation time	[23]
Transmissible Gastroenteritis Virus (TGEV)	254	3.68	liquid (with 10% blood platelet concentrate)	probably not highly absorbed, values taken from Figure 3A	[11]
Transmissible Gastroenteritis Virus (TGEV)	254	11.36	liquid (with 30% blood platelet concentrate)	probably highly absorbed, values taken from Figure 3A	[11]
Canine Coronavirus (CCoV S378)	254	1473*	liquid (cell culture medium with fetal bovine serum in MTP)	probably highly absorbed, UV lamp 4 cm away from MTP, sample evaporation? – not included in analysis	[12]
Human Coronavirus (HCoV-229E)	254	successful inactivation	liquid (cell culture medium with serum)	probably highly absorbed, no quantification possible	[25]
Human Coronavirus (HCoV-OC43)	254	successful inactivation	liquid (cell culture medium with serum)	probably highly absorbed, no quantification possible	[25]

(Continued)

Table 1: Overview of published and analyzed coronavirus photoinactivation investigations sorted by sample condition

Infectious Bronchitis Corona Virus (IBV-10 strains)	254	successful inactivation	liquid (cell culture medium)	no dose information, "no marked differences between strains"	[26]
Middle East Respiratory Syndrome Coronavirus (MERS-CoV EMC 2012)	254	27.47	liquid (blood platelet concentrate)	probably highly absorbed	[27]
Murine Coronavirus (MHV)	254	6.67	liquid (cell culture medium without serum)	probably highly absorbed, value estimated by statements in the Materials and methods section	[28]
Porcine Epidemic Diarrhea Virus (PEDV)	254	successful inactivation	liquid (blood plasma)	high absorption, no quantification possible	[29]
Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV Frankfurt 1)	254	17.54	liquid (blood platelet concentrate)	probably highly absorbed	[30]
Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV Hanoi)	254	22.67	liquid (cell culture medium with fetal bovine serum in MTP)	probably highly absorbed	[31]
Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV P9)	254	40.5	liquid (cell culture medium with fetal bovine serum in MTP)	probably highly absorbed, values estimated from semi-quantitative data in Table 3	[32]
Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV Urbani)	254	321*	liquid (cell culture medium with fetal bovine serum in MTP)	probably highly absorbed, UV lamp 3 cm away from MTP, sample evaporation?, values taken from Figure 1(A) – not included in analysis	[13]
Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV Urbani)	254	1135*	liquid (salt solution in MTP)	UV lamp 3 cm away from MTP, sample evaporation?, values taken from Figure 2 – not included in analysis	[10]
Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV Urbani)	254	11754*	liquid (salt solution with bovine serum albumin)	probably highly absorbed, UV lamp 3 cm away from MTP, sample evaporation?, values taken from Figure 2 – not included in analysis	[10]
Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV Urbani)	365	4800*	liquid (cell culture medium with fetal bovine serum in MTP)	UV lamp 3 cm away from MTP, sample evaporation?, values taken from Figure 1(A), no UVC – not included in analysis	[13]
Bovine Coronavirus (BCoV)	254	0.6*	no information available	data not included in analysis	[9]
Canine Coronavirus (CCoV)	254	0.6*	no information available	data not included in analysis	[9]
Infectious Bronchitis Corona Virus (IBV)	254	23*	no information available	data not included in analysis	[9]
Porcine Respiratory Coronavirus (PRCV)	254	1.3*	no information available	data not included in analysis	[9]

Values marked by * are not included in the further analysis, for reasons explained in the "Results and discussion" section. The log-reduction doses give the declared or determined necessary irradiation dose for a 90% virus inactivation.

2. Several investigators performed their irradiation experiments in microtiter plates (MTPs). This might become a problem if the plates are too close to the irradiation source, e.g., in the range of only a few centimeters. Besides the risk of heating the sample and subsequently increased sample evaporation, determination of the irradiation intensity inside the MTP wells becomes difficult. The MTP well walls shade the virus-containing samples from irradiation that does not originate directly above the well. Hence, the true irradiation intensity within a well is probably quite low compared to the intensity measured by a photodetector at the same distance.

Table 1 states whether a study exhibited one or two of these complications for the intended log-reduction determination. Actually, all calculated extreme values in Table 1 seem to be influenced by both complicating factors, and therefore these 254 nm values [12], [13], [10] were omitted in the further analysis. The data of the Berne torovirus [14] were also excluded, because of structural differences between toroviruses and coronaviruses [15].

All coronaviruses exhibit a similar structure and a single-stranded RNA length of about 30 kb, allowing the conclusion that they also feature very similar UVC absorption and UVC disinfection properties. It is therefore justified to consider all coronaviruses alike in terms of the investigated UVC-based log-reduction and to summarize the individual results.

This leads to a total UVC median log-reduction dose of 10.6 mJ/cm² (average 11.9±11.4 mJ/cm²). These values were calculated without the torovirus data and outliers, but the input included viruses in media that probably had higher UVC absorption, leading to reduced photoinactivation. Therefore, this 10.6 mJ/cm² is probably not the real value for the log-reduction dose, but instead could be considered as an upper limit.

Recalculation of the UVC log-reduction dose without using results from higher-absorption media should lead to more realistic values. In this case, the total median log-reduction dose would be 3.7 mJ/cm² (average 5.8±5.5 mJ/cm²).

This overview covers all coronaviruses and both UVC wavelengths (222 nm and 254 nm), but without results obtained from studies with probably high absorption media. The obtained results agree well with UVC inactivation data for other ssRNA viruses, such as influenza A with log-reduction doses around 2 mJ/cm² [5] or the ssRNA bacteriophage MS2 with log-reduction doses of about 20 mJ/cm² [5].

So far, most instances of successful coronavirus inactivation have been performed using mercury vapor lamps with peak emission at 254 nm. To reduce the use of the toxic mercury, it seems possible that these vapor lamps will be replaced in the future by 222-nm excimer lamps or by 270-nm LED. Since the RNA absorption strengths are similar, the disinfecting effect at these wavelengths will probably be approximately the same as with mercury

vapor lamps. However, this should be investigated in more detail in the future, since absorptions in the lipid envelope might have a larger influence on virus inactivation than currently assumed.

Because there are only single results available for the effect of 365 nm (UVA) and daylight, the focus of this analysis is on UVC coronavirus inactivation. Both irradiation methods demonstrated a virus reduction, albeit seemingly much less effective than that achieved with 254 nm irradiation. Nevertheless, these longer wavelengths might also be of future interest because their absorption in samples with organic materials is much lower, resulting in higher penetration depths which may allow virus inactivation of larger volumes.

Conclusion

To date, UVC radiation has been effective against all coronaviruses in all published investigations, although the absorption properties of the sample media reduced inactivation success. The calculated upper limit for the log-reduction median dose (in low-absorbance media) is 10.6 mJ/cm², but the probably more precise estimation is 3.7 mJ/cm².

These results were obtained by investigations on many different coronaviruses, including SARS-CoV and MERS-CoV, but not SARS-CoV-2. Nevertheless, it can be assumed that they are also applicable for SARS-CoV-2 and all future mutations. RNA mutations might have a strong influence on the pathogenicity of a virus, but they do not result in larger structural differences, especially concerning the UV absorption properties of the RNA, which are the main cause for the antiviral effect of ultraviolet radiation.

The above-mentioned log-reduction doses are in the same order of magnitude or even smaller than log-reduction doses for other important pathogens, such as *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumoniae* or *Candida albicans* [5]. They are also low compared to UVC irradiation recommendations, for instance, the international standard for UV disinfection of drinking water [16] with its recommended irradiation dose of 40 mJ/cm². Therefore, it can be concluded that existing UVC disinfection systems and procedures will be sufficient to deal with all coronaviruses, including SARS-CoV-2.

Notes

Competing interests

The authors declare that they have no competing interests.

References

- Johns Hopkins University. Coronavirus Resource Center – COVID-19 Dashboard by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University. [accessed 2020 May 6]. Available from: <https://coronavirus.jhu.edu/map.html>
- Li X, Luk HKH, Lau SKP, Woo PCY. Human Coronaviruses: General Features. In: Reference Module in Biomedical Sciences. Amsterdam: Elsevier; 2019. p. 1-6. DOI: 10.1016/B978-0-12-801238-3.95704-0
- Chang L, Yan Y, Wang L. Coronavirus Disease 2019: Coronaviruses and Blood Safety. *Transfus Med Rev.* 2020 Feb 21. pii: S0887-7963(20)30014-6. DOI: 10.1016/j.tmr.2020.02.003
- Kampf G, Voss A, Scheithauer S. Inactivation of coronaviruses by heat. *J Hosp Infect.* 2020 Mar 31. DOI: 10.1016/j.jhin.2020.03.025
- Kowalski W. Ultraviolet Germicidal Irradiation Handbook. Berlin, Heidelberg: Springer; 2009.
- Jagger J. Introduction to Research in Ultraviolet Photobiology. *Photochem Photobiol.* 1968;7:413. DOI: 10.1111/j.1751-1097.1968.tb08029.x
- Budowsky EI, Bresler SE, Friedman EA, Zheleznova NV. Principles of selective inactivation of viral genome. I. UV-induced inactivation of influenza virus. *Arch Virol.* 1981;68(3-4):239-47. DOI: 10.1007/BF01314577
- Chevrefils G, Caron É. UV dose required to achieve incremental Log inactivation of bacteria, protozoa and viruses. *IUVA News.* 2006;8(1):38-45.
- Kowalski W. Performance of the UV24 Unit Against Zoonotic Pathogens. San Fernando, CA: Medical Illumination. 2017 Feb 1. Available from: <http://www.medillum.com/wp-content/uploads/2017/03/Performance-of-the-UV24-Unit-Against-Zoonotic-Pathogens.pdf>
- Darnell MER, Taylor DR. Evaluation of inactivation methods for severe acute respiratory syndrome coronavirus in noncellular blood products. *Transfusion.* 2006 Oct;46(10):1770-7. DOI: 10.1111/j.1537-2995.2006.00976.x
- Terpstra FG, van 't Wout AB, Schuitemaker H, van Engelenburg FAC, Dekkers DWC, Verhaar R, de Korte D, Verhoeven AJ. Potential and limitation of UVC irradiation for the inactivation of pathogens in platelet concentrates. *Transfusion.* 2008 Feb;48(2):304-13. DOI: 10.1111/j.1537-2995.2007.01524.x
- Pratelli A. Canine coronavirus inactivation with physical and chemical agents. *Vet J.* 2008 Jul;177(1):71-9. DOI: 10.1016/j.tvjl.2007.03.019
- Darnell MER, Subbarao K, Feinstone SM, Taylor DR. Inactivation of the coronavirus that induces severe acute respiratory syndrome. SARS-CoV. *J Virol Methods.* 2004 Oct;121(1):85-91. DOI: 10.1016/j.jviromet.2004.06.006
- Weiss M, Horzinek MC. Resistance of Berne virus to physical and chemical treatment. *Vet Microbiol.* 1986 Feb;11(1):41-9. DOI: 10.1016/0378-1135(86)90005-2
- Cavanagh D. Coronaviruses and Toroviruses. In: Zuckerman AJ, Banatvala JE, Pattison JR, editors. Principles and practice of clinical virology. 4th ed. Chichester: Wiley; 2000. p. 345-56. DOI: 10.1002/0470842474.ch10
- DIN Deutsches Institut für Normung e.V. DIN EN 14897:2007-09 – Anlagen zur Behandlung von Trinkwasser innerhalb von Gebäuden – Geräte mit Quecksilberdampf-Niederdruckstrahlern – Anforderungen an Ausführung, Sicherheit und Prüfung [DIN EN 14897:2007-09 – Water conditioning equipment inside buildings – Devices using mercury low-pressure ultraviolet radiators – Requirements for performance, safety and testing]. Berlin: Beuth; 2007.
- Buonanno M, Welch D, Shuryak I, Brenner DJ. Far-UVC light efficiently and safely inactivates airborne human coronaviruses. (Preprint, Version 1). *Research Square.* 2020 Apr 27. DOI: 10.21203/rs.3.rs-25728/v1
- Walker CM, Ko G. Effect of ultraviolet germicidal irradiation on viral aerosols. *Environ Sci Technol.* 2007 Aug 1;41(15):5460-5. DOI: 10.1021/es070056u
- Bedell K, Buchaklian AH, Perlman S. Efficacy of an Automated Multiple Emitter Whole-Room Ultraviolet-C Disinfection System Against Coronaviruses MHV and MERS-CoV. *Infect Control Hosp Epidemiol.* 2016 May;37(5):598-9. DOI: 10.1017/ice.2015.348
- Deshmukh DR, Pomeroy BS. Ultraviolet inactivation and photoreactivation of avian viruses. *Avian Dis.* 1969 Aug;13(3):596-602.
- Cartwright SF. A cytopathic virus causing a transmissible gastroenteritis in swine. II. Biological and serological studies. *J Comp Pathol.* 1966 Jan;76(1):95-106. DOI: 10.1016/0021-9975(66)90051-x
- Ansaldi F, Banfi F, Morelli P, Valle L, Durando P, Sticchi L, Contos S, Gasparini R, Crovari P. SARS-CoV, influenza A and syncytial respiratory virus resistance against common disinfectants and ultraviolet irradiation. *J Prev Med Hyg.* 2004 Mar;45(1-2):5-8.
- Saknimit M, Inatsuki I, Sugiyama Y, Yagami K. Virucidal efficacy of physico-chemical treatments against coronaviruses and parvoviruses of laboratory animals. *Jikken Dobutsu.* 1988 Jul;37(3):341-5. DOI: 10.1538/expanim1978.37.3_341
- Hirano N, Hino S, Fujiwara K. Physico-chemical properties of mouse hepatitis virus (MHV-2) grown on DBT cell culture. *Microbiol Immunol.* 1978;22:377-90. DOI: 10.1111/j.1348-0421.1978.tb00384.x
- Bucknall RA, King LM, Kapikian AZ, Chanock RM. Studies with human coronaviruses. II. Some properties of strains 229E and OC43. *Proc Soc Exp Biol Med.* 1972;139(3):722-7. DOI: 10.3181/00379727-139-36224
- Otsuki K, Yamamoto H, Tsubokura M. Studies on avian infectious bronchitis virus (IBV): I. Resistance of IBV to chemical and physical treatments. *Arch Virol.* 1979;60(1):25-32. DOI: 10.1007/BF01318094
- Eickmann M, Gravemann U, Handke W, Tolksdorf F, Reichenberg S, Müller TH, Seltsam A. Inactivation of Ebola virus and Middle East respiratory syndrome coronavirus in platelet concentrates and plasma by ultraviolet C light and methylene blue plus visible light, respectively. *Transfusion.* 2018 Sep;58(9):2202-7. DOI: 10.1111/trf.14652
- Liu Y, Cai Y, Zhang X. Induction of caspase-dependent apoptosis in cultured rat oligodendrocytes by murine coronavirus is mediated during cell entry and does not require virus replication. *J Virol.* 2003 Nov;77(22):11952-63. DOI: 10.1128/jvi.77.22.11952-11963.2003
- Blázquez E, Rodríguez C, Ródenas J, Navarro N, Riquelme C, Rosell R, Campbell J, Crenshaw J, Segalés J, Pujols J, Polo J. Evaluation of the effectiveness of the SurePure Turbulator ultraviolet-C irradiation equipment on inactivation of different enveloped and non-enveloped viruses inoculated in commercially collected liquid animal plasma. *PLoS ONE.* 2019 Feb 21;14(2):e0212332. DOI: 10.1371/journal.pone.0212332

30. Eickmann M, Gravemann U, Handke W, Tolksdorf F, Reichenberg S, Müller TH, Seltsam A. Inactivation of three emerging viruses – severe acute respiratory syndrome coronavirus, Crimean-Congo haemorrhagic fever virus and Nipah virus – in platelet concentrates by ultraviolet C light and in plasma by methylene blue plus visible light. *Vox Sang*. 2020 Apr;115(3):146-51. DOI: 10.1111/vox.12888
31. Kariwa H, Fujii N, Takashima I. Inactivation of SARS coronavirus by means of povidone-iodine, physical conditions, and chemical reagents. *Jpn J Vet Res*. 2004 Nov;52(3):105-12.
32. Duan SM, Zhao XS, Wen RF, Huang JJ, Pi GH, Zhang SX, Han J, Bi SL, Ruan L, Dong XP; SARS Research Team. Stability of SARS coronavirus in human specimens and environment and its sensitivity to heating and UV irradiation. *Biomed Environ Sci*. 2003 Sep;16(3):246-55.

Corresponding author:

Prof. Dr. Martin Heßling
Institute of Medical Engineering and Mechatronics, Ulm
University of Applied Sciences, Albert-Einstein-Allee 55,
89081 Ulm, Germany
Martin.Hessling@thu.de

Please cite as

Heßling M, Hönes K, Vatter P, Lingensfelder C. Ultraviolet irradiation doses for coronavirus inactivation – review and analysis of coronavirus photoinactivation studies. *GMS Hyg Infect Control*. 2020;15:Doc08. DOI: 10.3205/dgkh000343, URN: urn:nbn:de:0183-dgkh0003436

This article is freely available from

<https://www.egms.de/en/journals/dgkh/2020-15/dgkh000343.shtml>

Published: 2020-05-14

Copyright

©2020 Heßling et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License. See license information at <http://creativecommons.org/licenses/by/4.0/>.